

Pollutant particles produce vasoconstriction and enhance MAPK signaling via angiotensin type I receptor

Zhuowei Li, Jacqueline D. Carter, Lisa A. Dailey, and
Yuh-Chin T. Huang

doi:10.1289/ehp.7736 (available at <http://dx.doi.org/>)
Online 15 April 2005



**Pollutant particles produce vasoconstriction and enhance MAPK signaling via
angiotensin type I receptor**

Zhuowei Li¹, Jacqueline D. Carter², Lisa A. Dailey², Yuh-Chin T. Huang²

¹Center for Environmental Medicine, Asthma and Lung Biology

University of North Carolina, Chapel Hill, NC 27599

and

²National Health and Environmental Effects Research Laboratory

Office of Research and Development

Environmental Protection Agency, Research Triangle Park, NC 27711

Correspondence: Yuh-Chin T. Huang, M.D., M.H.S.

CB 7315, 104 Mason Farm Road

Chapel Hill, NC 27599

919-843-9504

FAX: 919-966-6271

E-mail: huang.tony@epa.gov

Running title: Particles and angiotensin

Keywords: air pollutant; angiotensin II; p38, ERK, vanadium, copper, angiotensin-converting enzyme

Acknowledgement: The authors would like to thank Craig Marshall of Duke University Medical Center for procuring rat pulmonary artery rings. Portion of the data were presented in the International Meeting of the American Thoracic Society in 2004 (Am. J. Respir. Crit. Care Med. 169: A168, 2004)

Abbreviations: angiotensin type 1 receptor (AT₁R), particulate matter (PM), human pulmonary artery endothelial cells (HPAECs), nitric oxide (NO), epidermal growth factor receptor (EGFR), angiotensin converting enzyme (ACE), mitogen-activated protein kinase (MAPK), extracellular signal-regulated kinases ½ (ERK1/2).

ABSTRACT

INTRODUCTION

MATERIALS AND METHODS

RESULTS

DISCUSSION

CONCLUSIONS

DISCLAIMER

REFERENCES

FIGURE LEGENDS

FIGURES

ABSTRACT

Exposure to particulate matter (PM) is associated with acute cardiovascular mortality and morbidity, but the mechanisms are not entirely clear. In this study, we hypothesized that PM may activate the angiotensin type 1 receptor (AT₁R), a G protein-coupled receptor that regulates inflammation and vascular function. We investigated the acute effects of St. Louis urban particle (UP, SRM1648) on the constriction of isolated rat pulmonary artery rings and the activation of extracellular signal-regulated kinases1/2 (ERK1/2) and p38 mitogen-activated protein kinases (MAPKs) in human pulmonary artery endothelial cells (HPAECs) with or without losartan, an antagonist of AT₁R. UP at 1-100 µg/ml induced acute vasoconstriction in pulmonary artery. UP also produced a time- and dose-dependent increase in phosphorylation of ERK1/2 and p38 MAPK. Losartan pretreatment inhibited both the vasoconstriction and the activation of ERK1/2 and p38. The water-soluble fraction of UP was sufficient for inducing ERK1/2 and p38 phosphorylation, which was also losartan-inhibitable. Copper and vanadium, two soluble transition metals contained in UP, induced pulmonary vasoconstriction and phosphorylation of ERK1/2 and p38, but only the phosphorylation of p38 was inhibited by losartan. The UP-induced activation of ERK1/2 and p38 was attenuated by captopril, an angiotensin-converting enzyme inhibitor. These results indicate that activation of the local renin-angiotensin system may play an important role in cardiovascular effects induced by PM.